

IA Medicaid Member ID # _ _ _ _ _ _ _ _ _ _ _ _ _ _	Patient name	DOB
Patient address		
Provider NPI _ _ _ _ _ _ _ _ _ _ _ _ _ _	Prescriber name	Phone
Prescriber address		Fax
Pharmacy name	Address	Phone
Prescriber must complete all information above. It must be legible, correct, and complete or form will be returned.		
Pharmacy NPI _ _ _ _ _ _ _ _ _ _ _ _ _ _	Pharmacy fax	NDC _ _ _ _ _ _ _ _ _ _ _ _ _ _

1. Patient is not using or planning to use a JAK inhibitor in combination with other JAK inhibitors, biological therapies, or potent immunosuppressants (azathioprine or cyclosporine); and
2. Request adheres to all FDA approved labeling for requested drug and indication, including age, dosing, contraindications, warnings and precautions, drug interactions, and use in specific populations; and
3. Patient has a diagnosis of:
 - a. Moderate to severe rheumatoid arthritis (baricitinib, tofacitinib, upadacitinib); with
 - i. A documented trial and inadequate response, at a maximally tolerated dose, with methotrexate; and
 - ii. A documented trial and inadequate response to one preferred TNF inhibitor; or
 - b. Psoriatic arthritis (tofacitinib, upadacitinib); with
 - i. A documented trial and inadequate response, at a maximally tolerated dose, with methotrexate (leflunomide or sulfasalazine may be used if methotrexate is contraindicated); and
 - ii. Documented trial and therapy failure with one preferred TNF inhibitor used for psoriatic arthritis; or
 - c. Moderately to severely active ulcerative colitis (tofacitinib, upadacitinib); with
 - i. A documented trial and inadequate response with a preferred TNF inhibitor; or
 - d. Moderately to severely active Crohn's disease (upadacitinib); with
 - i. A documented trial and inadequate response with a preferred TNF inhibitor; or
 - e. Polyarticular Course Juvenile Idiopathic Arthritis (tofacitinib); with
 - i. A documented trial and inadequate response to the preferred oral DMARD, methotrexate (leflunomide or sulfasalazine may be used if methotrexate is contraindicated); and
 - ii. A documented trial and inadequate response with a preferred TNF inhibitor; or
 - f. Axial spondyloarthritis conditions (e.g., ankylosing spondylitis or nonradiographic axial spondyloarthritis) (tofacitinib, upadacitinib); with
 - i. A documented trial and inadequate response to at least two preferred non-steroidal anti-inflammatories (NSAIDs) at a maximally tolerated dose for a minimum of at least one month; and
 - ii. A documented trial and inadequate response with at least one preferred TNF inhibitor; or
 - g. Atopic dermatitis; with
 - i. Documentation patient has failed to respond to good skin care and regular use of emollients; and
 - ii. A documented adequate trial and therapy failure with one preferred medium to high potency topical corticosteroid for a minimum of 2 consecutive weeks; or
 - iii. A documented trial and therapy failure with a topical immunomodulator for a minimum of 4 weeks; and
 - iv. For mild to moderate atopic dermatitis (topical treatments):
 - a. Affected area is less than 20% of body surface area (BSA); and
 - b. Patient has been instructed to use no more than 60 grams of topical ruxolitinib per week; or
 - v. For moderate to severe chronic hand eczema (topical treatments):
 - a. Chronic hand eczema has persisted for more than 3 months or recurred two or more times with a 12-month time frame after the initial occurrence with complete clearances between relapses; and
 - b. Patient has been instructed to use no more than 30 grams per 2 weeks or 60 grams per month of topical delgocitinib; or

Request for Prior Authorization
JANUS KINASE (JAK) INHIBITORS
(PLEASE PRINT – ACCURACY IS IMPORTANT)

vi. For moderate to severe atopic dermatitis (oral treatments):

- a. A documented trial and therapy failure with a systemic drug product for the treatment of moderate to severe atopic dermatitis, including biologics; and
- b. Requests for upadacitinib for pediatric patients 12 to less than 18 years if age must include the patient's weight in kg; or
- h. Nonsegmental vitiligo (ruxolitinib) with;
 - i. A documented trial and inadequate response with a potent topical corticosteroid; or
 - ii. A documented trial and inadequate response with a topical calcineurin inhibitor; and
 - iii. The patient's body surface area (BSA) is less than or equal to the affected BSA per FDA approved label, if applicable; or
- i. Giant Cell Arteritis; with
 - i. Documentation patient is currently taking a glucocorticoid, with a tapering dose, or has discontinued use of glucocorticoids.

The required trials may be overridden when documented evidence is provided that use of these agents would be medically contraindicated.

Preferred

☐ Opzelura ☐ Xeljanz

☐ Rinvoq

Non-Preferred

☐ Anzupgo ☐ Olumiant ☐ Xeljanz XR

☐ Cibinqo ☐ Xeljanz Oral Solution

Strength _____ **Dosage Instructions** _____ **Quantity** _____ **Days Supply** _____

Diagnosis: _____

Will the JAK inhibitor be used in combination with other JAK inhibitors, biological therapies or potent immunosuppressants? ☐ Yes ☐ No

☐ **Moderate to Severe Rheumatoid Arthritis (RA) (Olumiant, Rinvoq, Xeljanz or Xeljanz XR)**

Methotrexate trial: Dose: _____ Trial dates: _____

Failure reason: _____

Preferred TNF Inhibitor: Name/Dose: _____ Trial Dates: _____

Failure reason: _____

☐ **Psoriatic Arthritis (Rinvoq, Xeljanz or Xeljanz XR)**

Methotrexate trial (leflunomide or sulfasalazine if methotrexate is contraindicated):

Name/Dose: _____ Trial dates: _____

Failure reason: _____

Preferred TNF Inhibitor: Name/Dose: _____ Trial Dates: _____

Failure reason: _____

☐ **Ulcerative Colitis (Rinvoq, Xeljanz or Xeljanz XR)**

Preferred TNF Inhibitor: Name/Dose: _____ Trial Dates: _____

Failure reason: _____

If requesting continuation of tofacitinib 10mg twice daily dose, document adequate therapeutic benefit:

☐ **Moderately to severely active Crohn's disease (Rinvoq)**

Preferred TNF Inhibitor: Name/Dose: _____ Trial Dates: _____

Failure reason: _____

Request for Prior Authorization
JANUS KINASE (JAK) INHIBITORS
(PLEASE PRINT – ACCURACY IS IMPORTANT)

☐ **Polyarticular Course Juvenile Idiopathic Arthritis (Xeljanz)**

Methotrexate trial (leflunomide or sulfasalazine if methotrexate is contraindicated):

Name/Dose: _____ Trial dates: _____

Failure reason: _____

Preferred TNF Inhibitor: Name/Dose: _____ Trial Dates: _____

Failure reason: _____

☐ **Axial spondyloarthritis conditions (e.g., ankylosing spondylitis or nonradiographic axial spondyloarthritis) (Rinvoq, Xeljanz or Xeljanz XR)**

Preferred NSAID trial 1: Name/Dose: _____ Trial Dates: _____

Failure reason: _____

Preferred NSAID trial 2: Name/Dose: _____ Trial dates: _____

Failure reason: _____

Preferred TNF Inhibitor: Name/Dose: _____ Trial Dates: _____

Failure reason: _____

☐ **Atopic Dermatitis**

Has patient failed to respond to good skin care and regular use of emollients? ☐ Yes ☐ No

Document emollient use: Product name, dosing instructions & duration of use: _____

Document trial and therapy failure with one preferred medium to high potency topical corticosteroid for a minimum of 2 weeks or topical immunomodulator for a minimum of 4 weeks:

Preferred Medium to High Potency Topical Corticosteroid Trial:

Drug name & dose: _____ Trial dates: _____

Failure reason: _____

Preferred Topical Immunomodulator Trial:

Drug name & dose: _____ Trial dates: _____

Failure reason: _____

Mild to Moderate Atopic Dermatitis (Topical Treatments)

Is affected area less than 20% of body surface area? ☐ Yes ☐ No

Has patient been instructed to use no more than 60gms of topical ruxolitinib per week? ☐ Yes ☐ No

Moderate to Severe Chronic Hand Eczema (Topical Treatments)

Has hand eczema persisted for more than 3 months or recurred two or more times within a 12-month timeframe after the initial occurrence with complete clearance between relapses? ☐ Yes ☐ No

Has patient been instructed to use no more than 30gms per 2 weeks or 60gms per month of topical delgocitinib?

☐ Yes ☐ No

Moderate to Severe Atopic Dermatitis (Oral Treatments)

Trial with systemic drug product for the treatment of moderate to severe atopic dermatitis, including biologics:

Drug name & dose: _____ Trial dates: _____

Failure reason: _____

Request for Prior Authorization
JANUS KINASE (JAK) INHIBITORS
(PLEASE PRINT – ACCURACY IS IMPORTANT)

Requests for upadacitinib for pediatric patients 12 to less than 18 years of age include weight in kg:

☐ **Nonsegmental vitiligo (Opzelura)**

Potent Topical Corticosteroid Trial:

Drug name & dose: _____ Trial dates: _____

Failure reason: _____

Topical Calcineurin Inhibitor Trial:

Drug name & dose: _____ Trial dates: _____

Failure reason: _____

Provide patient's affected body surface area (BSA): _____

☐ **Giant Cell Arteritis**

Is patient currently taking a glucocorticoid?

☐ Yes; Is dose being tapered? ☐ Yes ☐ No

☐ No

Other medical conditions to consider: _____

Attach lab results and other documentation as necessary.

Prescriber signature (Must match prescriber listed above.)	Date of submission
--	--------------------

IMPORTANT NOTE: In evaluating requests for prior authorization the consultant will consider the treatment from the standpoint of medical necessity only. If approval of this request is granted, this does not indicate that the member continues to be eligible for Medicaid. It is the responsibility of the provider who initiates the request for prior authorization to establish by inspection of the member's Medicaid eligibility card and, if necessary, by contact with the county Department of Health and Human Services, that the member continues to be eligible for Medicaid.